



The acceptability, feasibility, and possible benefits of a neurobiologically-informed 5-day multifamily treatment for adults with anorexia nervosa

Christina E. Wierenga PhD¹  | Laura Hill PhD² | Stephanie Knatz Peck PhD¹ | Jason McCray PhD² | Laura Greathouse MPH¹ | Danika Peterson BA¹ | Amber Scott BA² | Ivan Eisler PhD³ | Walter H. Kaye MD¹ 

¹Department of Psychiatry, University of California San Diego, San Diego, California

²The Center for Balanced Living, Columbus, Ohio

³South London and Maudsley NHS Foundation Trust, London, United Kingdom

Correspondence

Christina E. Wierenga, PhD, Associate Professor of Psychiatry, UCSD Department of Psychiatry, Chancellor Park, 4510 Executive Dr., Suite 315, San Diego, CA 92121.

Email: cwierenga@ucsd.edu; <http://eatingdisorders.ucsd.edu>

Funding information

National Eating Disorders Association (NEDA); the Global Foundation for Eating Disorders (GFED) and the Price Foundation

Abstract

Objective: Novel treatments for adults with anorexia nervosa (AN) are lacking. Recent scientific advances have identified neurobiologically-driven temperament contributors to AN symptoms that may guide development of more effective treatments. This preliminary study evaluates the acceptability, feasibility and possible benefits of a multicenter open trial of an intensive 5-day neurobiologically-informed multifamily treatment for adults with AN and their supports (SU). The temperament-focused treatment combines psychoeducation of AN neurobiology and SU involvement to develop skills to manage traits contributing to disease chronicity.

Method: Fifty-four adults with AN and at least one SU ($n = 73$) received the 5-day treatment. Acceptability, feasibility, and attrition were measured post-treatment. Clinical outcome (BMI, eating disorder psychopathology, family function) was assessed post-treatment and at >3-month follow-up.

Results: The treatment had low attrition, with only one drop-out. Patients and SU rated the intervention as highly acceptable, and clinicians reported good feasibility. At post-treatment, patients demonstrated significantly increased BMI, reduced eating disorder psychopathology, and improved family function. Benefits were maintained in the 39 patients who completed follow-up assessment, with 62% reporting full or partial remission.

Discussion: Preliminary results are promising and suggest this novel treatment is feasible and acceptable. To establish treatment efficacy, fully-powered randomized controlled trials are necessary.

KEYWORDS

anorexia nervosa, intensive multifamily treatment, open trial, temperament based treatment

1 | INTRODUCTION

Anorexia nervosa (AN) is a serious psychiatric disorder that is difficult to treat and is associated with significant medical and psychological complications. Treatment efficacy in adults is limited, with no treatment showing superiority (NICE, 2017). Approximately 50% of patients develop a persistent, relapsing, and remitting course (Hay, Touyz, & Sud, 2012; Steinhausen, 2002). As such, identifying efficacious treatments for adults with AN is imperative (Hay et al., 2012; Watson & Bulik, 2013). Unfortunately, the etiology of AN is poorly

understood, hindering the ability of treatments to target underlying mechanisms.

Accumulating behavioral and neuroimaging evidence supports a neurobiologically-based temperament that impacts the development and maintenance of AN—characterized by anxiety, reward insensitivity, perfectionism, altered interoceptive awareness, harm avoidance, and cognitive inflexibility (Fassino, Piero, Gramaglia & Abbate-Daga, 2004; Harrison, O'Brien, Lopez, & Treasure, 2010; Lilenfeld, 2011)—that is related to altered insula and fronto-striatal neural circuit function (Berner et al., 2017; DeGuzman, Shott, Yang, Riederer, & Frank, 2017;

Kerr et al., 2016; Oberndorfer et al., 2013; Wierenga et al., 2015). In addition to preexisting AN, mild to modest amounts of these traits often persist after recovery (Wagner et al., 2006), suggesting those who recover might do so by effectively managing these traits. This personality and behavior profile establishes a framework to guide development of therapies designed to directly target trait-related symptoms specific to AN. Yet, to date only a few therapies, such as Maudsley model of anorexia nervosa treatment for adults (MANTRA) (Schmidt et al., 2013), are based on such an empirically supported understanding of AN, and none are administered in an intensive multifamily format with neurobiological psychoeducation and skills training.

To address this need and respond to the recent call that treatments for AN not remain “brainless” (Schmidt & Campbell, 2013), we developed an intensive 5-day multifamily treatment for adult AN (formerly called intensive temperament based therapy and neurobiologically enhanced with family eating disorder trait response treatment [NEWFED-TR]) (Kaye et al., 2015; Knatz, Wierenga, Murray, Hill, & Kaye, 2015). The treatment includes psychoeducation to emphasize the key role of neurobiological factors in the development of AN, teaching the patient and Support (SU) age-appropriate skills to manage disorder-related temperament (see Supporting Information for details, including the schedule of treatment modules and activities). We previously demonstrated clinical effects following a 1-week intervention for adolescent AN (Marzola et al., 2015; Rockwell, Boutelle, Trunko, Jacobs, & Kaye, 2011). This intensive format was adapted for adults while incorporating different interventions specific to adult AN (Hill, 2017; Hill, Peck, Wierenga, & Kaye, 2016) (see Supporting Information). The 5-day treatment leverages SU participation based on considerable data showing that AN-focused family therapy (FT-AN) is the most effective approach in treating adolescents with AN (Eisler, Grange, & Lock, 2015; Lock, 2015; NICE, 2017), with multifamily therapy showing improved efficacy over single-family therapy (Eisler et al., 2016). In adult AN, family therapy has evidenced superiority to treatment as usual and efficacy equal to other specialist eating disorder (ED) therapies (Dare, Eisler, Russell, Treasure, & Dodge, 2001), with considerable evidence that family interventions in adult mental health can be enhanced by using a multifamily treatment format (Lemmens, Eisler, Buysse, Heene, & Demyttenaere, 2009; McFarlane, 2016; Miller, Solomon, Ryan, & Keitner, 2004), and is feasible in adult AN (Dimitropoulos, Farquhar, Freeman, Colton, & Olmsted, 2015; Tantillo, 2006).

To our knowledge, an intensive (e.g., ~40 h) multifamily treatment format has not been implemented for adult AN. Thus, this preliminary study evaluated the feasibility and acceptability of an intensive, multifamily, temperament-focused treatment for adult AN, and secondarily explored changes in clinical outcome (e.g., self-reported ED symptoms, family function).

2 | METHODS

2.1 | Participants

Participants were recruited via advertisements on our clinic websites. Sixty-two adults were assessed for eligibility, of which seven met

DSM-IV-TR (American Psychiatric Association, 2000) criteria for bulimia nervosa (BN) or subthreshold BN and were excluded (see Supporting Information for additional exclusionary criteria). Fifty-five adult women met criteria for broadly defined AN (e.g., including DSM-IV-TR AN, AN in partial remission, and Eating Disorder Not Otherwise Specified, restricting type) as determined by consensus diagnosis and based on Module H of the Structured Clinical Interview for DSM-IV (First, Gibbon, Spitzer, & Williams, 1996) administered by trained clinicians. Patients selected their participating SU ($n = 73$): parents (36 mothers, 17 fathers), significant others (2 boyfriends, 2 fiancées, 5 husbands, 2 partners, 1 wife), other family members (4 sisters, 1 cousin, and 1 nanny), and friends (2). 39 patients were accompanied by 1 SU, 14 patients by 2 SU, and 2 patients by three SU.

The study conforms to the standards of the Declaration of Helsinki and the protocol was reviewed and approved by the Institutional Review Board of the University of California San Diego, which covered The Center for Balanced Living as a secondary site. All AN and SU participants provided written informed consent.

2.2 | Measures

2.2.1 | Feasibility

Qualitative feedback was gathered from study clinicians, patients, SU, and the multidisciplinary team regarding the program's practical implementation and the ease of adapting the intensive multifamily format to an adult population. This included an assessment of clinician willingness to recruit participants, staff time required, participant ability to engage with the activities, and estimation of follow-up rates.

2.2.2 | Acceptability

An 18-item Patient and SU Satisfaction Questionnaire developed by our team was used to assess patient and SU liking and any concerns related to the 5-day treatment (patient $\alpha = 0.92$; SU $\alpha = 0.88$). The measure also included an open-ended question to elicit qualitative feedback. Daily attendance was recorded and a daily feedback form was administered to participants to track dropout and evaluation of experiential activities.

2.2.3 | Clinical measures

Staff measured patients' height and gown weight immediately prior to and following 5-day treatment. Measurements were converted to body mass index (BMI; kg/m^2), and ideal body weight (%IBW) was calculated for sex and height using the 1959 Metropolitan Life Insurance table. Participants completed self-report assessments at entry, immediately post-treatment, and at follow-up (>3 months), including the Eating Disorder Examination Questionnaire (EDE-Q) (Fairburn & Beglin, 1994) with parent version (P-EDEQ; (Loeb, 2008) modified for SU, McMaster Family Assessment Device (FAD) (Epstein, Baldwin, & Bishop, 1983), and Spielberger State-Trait Anxiety Inventory (STAI) (Spielberger, Gorsuch, & Lushene, 1970). Cronbach's alphas for all measures within the present study were strong ($\alpha = .88-.97$). Secondary measures are reported in the Supporting Information.

2.3 | Exploratory outcome classification

Patients who completed assessments at pretreatment, post-treatment, and follow-up were classified as fully remitted, partially remitted, or poor outcome according to criteria set forth by (Bardone-Cone et al., 2016) (see Supporting Information for details).

2.4 | Statistical analysis

Parametric and nonparametric tests were selected as appropriate. Measures with missing values were excluded from analysis; missing values occurred on <1% of all self-reported items. Within-subject changes in clinical measures between pre- and post-treatment were analyzed using paired sample *t* tests. These analyses were repeated for SU responses, weighted for number of SU per patient. McNemar's exact tests were used to compare good (partial or full remission) vs. poor outcome classification of patients between pretreatment, post-treatment, and follow-up. Lastly, we examined whether treatment location predicted outcome and follow-up assessment completion using linear and logistic regression analyses (see Supporting Information for details of additional statistical analysis of follow-up data).

3 | RESULTS

3.1 | Feasibility

At each site, the treatment included an intake coordinator, three clinicians, a dietician, a physician, and a clinical administrative assistant. Including preparation, actual sessions, communicating with other treatment providers, discharge planning, and note writing, 80 h of staff time were required. This was perceived as feasible by the two administering programs, considering that up to 6 patients' and 6–15 SUs attendance equated to 13 h of staff time per patient for the entire intervention.

3.2 | Attrition and acceptability

One patient necessitated a higher level of care after attending one day of the 5-day program, resulting in an attrition rate of 1.8%. Patient and SU acceptability ratings were strong across the 18 questions (Table 1), and 88% of patients and 95% of SU reported treatment met their expectations. Objective and qualitative feedback indicated that patients and SU particularly liked the neurobiology psychoeducation (97%), activities (98%), group format (100%), and behavioral contracting (89%). Of treatment completers ($n = 54$), 7% of patients ($n = 4$), and 8% of SU ($n = 6$) did not complete clinical outcome assessments at post-treatment. Of post-treatment assessment completers ($n = 50$), 22% of patients did not complete follow-up self-report assessments ($n = 11$) and an additional 24% of patients ($n = 12$) completed neither follow-up self-report nor weight assessments. The primary reason provided for not completing follow-up assessments was that patients were doing better and thought the questions would be triggering. Sixty-nine percent of SU did not complete follow-up assessments.

3.3 | Patient characteristics pre- and post- 5-day treatment

Table 2 includes patient characteristics of the 50 women (primarily Caucasian, 84.6%) who completed assessments pre- and post-treatment at the University of California, San Diego ($n = 17$) or at The Center for Balanced Living, Columbus OH ($n = 33$). Immediately after 5-day treatment, there was a significant increase in BMI ($p = .001$, $d = 0.10$, 95% CI = -0.68 – 0.46), though this likely does not represent a clinically meaningful change. Patients also reported significant reductions in EDE-Q Global score ($p = .014$, $d = 0.27$, 95% CI = -0.15 – 0.69), state anxiety ($p < .001$, $d = 0.68$, 95% CI = -2.87 – 4.09), and improvements in FAD General Family Function ($p = .005$, $d = 0.36$, 95% CI = 0.20 – 0.51) (Table 2); see Supporting Information for follow-up data results.

3.4 | Support ratings post 5-day treatment

Data from 67 SU who completed pre- and post-treatment assessments, weighted to reflect 50 unique patients, revealed significant improvements in observed ED symptoms of their loved one (EDE-Q Global score [observer modified], $p < .00$, $d = 0.72$, 95% CI = 0.30 – 1.08), and general family function (FAD, $p < .01$, $d = 0.54$, 95% CI = 0.43 – 0.64) (Supporting Information Table 2; see Supporting Information for follow-up data).

3.5 | Patient outcome classification

At post-treatment, 31% of patients were classified as fully remitted (10%, $n = 4$) or partially remitted (21%, $n = 8$), while 69% reported a poor outcome. At follow-up, 62% of patients achieved either full remission (31%, $n = 12$) or partial remission (31%, $n = 12$), while 38% ($n = 15$) reported a poor outcome (Supporting Information Table 3 and Figure 2). A significant difference was found for the proportion of individuals with good outcome (i.e., full or partial remission) at pretreatment versus follow-up ($p < .001$) and at post-treatment versus follow-up ($p = .004$), but not at pre- versus post-treatment ($p = .125$). No associations were found between treatment site and outcome or failure to complete follow-up assessments ($ps > .05$).

4 | DISCUSSION

This study represents the first step in evaluating the feasibility, acceptability, and possible benefits of an intensive 5-day neurobiologically-informed intervention for adult AN that enlists SU involvement through multifamily neurobiologically based psychoeducation and skills-training, aimed at managing traits that contribute to AN. Consistent with previous findings in a smaller sample (Hill, 2017), the treatment is feasible, has low attrition and high acceptability. Among treatment completers, findings indicate increased BMI, reduced ED symptoms, reduced state anxiety, and improved family function post-treatment. Exploratory results indicate that 62% of patients who completed follow-up assessments achieved full or partial remission (aka, good outcome) at follow-up. No differences in outcome were detected between treatment sites.

TABLE 1 Acceptability ratings post-treatment

	Pt (N = 45)		SU (N = 67)			
1. I would recommend the 5-day program to others	4.5		4.8			
2. I would prefer additional group treatment sessions or exercises	3.6		3.9			
3. I would be willing to participate in additional group treatment sessions or exercises	4.1		4.5			
4. I enjoyed the learning about the neurobiology of eating disorders through the group exercises (e.g., nondominant hand writing exercise, brain wave)	4.7		4.8			
5. The exercises on neurobiology improved my understanding about my eating disorder	4.8		4.8			
6. I enjoyed the activities for learning/practicing effective communication with my Support(s)/loved one	4.5		4.7			
7. I feel that my Support(s) are equipped with more/better tools for supporting me through recovery/I feel that I am equipped with more/better tools for supporting my loved one through recovery	4.6		4.7			
8. I feel that I am better able to communicate with my Support(s)/loved one about my/her eating disorder	4.5		4.6			
9. I enjoyed working on developing a contract/treatment plan with my Support(s)/loved one	3.8		4.6			
10. I am more confident about my Support(s)/my ability to support me/my loved one through recovery	4.4		4.6			
11. I feel that my Support(s)/my role in my/my loved one's treatment has been clarified	4.3		4.5			
12. My relationship with my Support(s)/loved one has improved as a result of this treatment	4.3		4.2			
13. I believe my experience from this treatment will be helpful in decreasing the likelihood that I/my loved one will engage in behaviors such as restricting, over-exercising, purging, etc	4.4		4.4			
14. I believe this treatment will be helpful in either decreasing my/my loved one's anxiety and/or other negative emotions or improving my/my loved one's ability to cope with these emotions	4.0		4.4			
15. I enjoyed interacting with other patients and their Supports in the group	4.7		4.7			
16. I learned skills and ideas from the other Supports and patients that I can now apply to myself/to working with my loved one in treatment	4.4		4.6			
17. I felt supported by the other group members	4.5		4.7			
18. I plan to continue to have my Support(s) involved in my treatment/I plan to continue my involvement in my loved one's treatment	4.5		4.8			
Mean (SD)	4.4 (.31)		4.6 (.26)			
	Yes (%)		Somewhat (%)		No (%)	
	Pt	SU	Pt	SU	Pt	SU
19. Did this treatment meet your expectations?	88.4	95.4	11.6	3.1	0	1.5

Note. Pt = patient; SU = support.

Acceptability ratings based on a 5-point Likert scale with higher ratings indicating greater acceptability for questions 1–18. Values associated with Question 19 indicate the percentage of respondents who felt that the treatment met expectations.

However, 22–46% of patients and 69% of SU failed to complete follow-up assessments, indicating suboptimal acceptability of ongoing study participation following treatment.

Future studies are needed to establish the efficacy and explore potential mechanisms of this neurobiologically-informed multifamily intervention. The neurobiological focus may have imparted benefits, as another trait-focused treatment (MANTRA) has shown efficacy (Byrne et al., 2017; Schmidt et al., 2013). The intensive nature of treatment may also have conferred benefits. Treatment models for anxiety indicate that intense, repeated, and focused *in vivo* practice is key to altering biologically-driven avoidance behaviors by maximizing learning

through massed practice and allowing close monitoring of compliance (Abramowitz, Foa, & Franklin, 2003; Gallo, Chan, Buzzell, Whitton, & Pincus, 2012; Storch et al., 2007). The contribution of other treatment components (e.g., multifamily format, behavioral contracting, dietary involvement) to outcome requires further study.

Despite these promising findings, this study has several limitations. These preliminary results are from an open trial with treatment-seeking patients who have active SU, resulting in a possible selection bias that may reflect particularly motivated AN (and/or SU) who are mild/moderately ill. As such, findings may not be representative of more resistant, severely underweight AN who lack social support, and clinical

TABLE 2 Patient clinical variables at pre- and post-5-day treatment

	Pretreatment (n = 50) Mean (SD)	Post-treatment (n = 50) Mean (SD)	p	Cohen's d	95% CI
Age (years)	24.5 (8.8)				
Illness duration (years)	9.1 (8.6)				
BMI (kg/m ²)	18.1 (2.1)	18.3 (2.0)	0.001	-0.10	-0.68-0.46
IBW (%)	83.1 (9.2)	84.1 (8.9)	0.001	-0.11	-2.66-2.36
EDE-Q Global Score	3.5 (1.5)	3.1 (1.5)	0.014	0.27	-0.15-0.69
FAD general family function	2.2 (.59)	2.0 (.53)	0.005	0.36	0.20-0.51
STAI state	56.2 (12.8)	47.8 (12.3)	<0.001	0.68	-2.87-4.09
STAI trait	55.3 (10.3)	53.4 (10.0)	0.26	0.19	-2.67-2.96
n (%)					
ED diagnosis					
AN-R	26 (52%)				
AN-BP	12 (24%)				
EDNOS-restricting type	2 (4%)				
AN-partial remission	10 (20%)				
Comorbid diagnoses					
Major depressive disorder	14 (28%)				
Generalized anxiety disorder	21 (42%)				
Panic disorder	7 (14%)				
Social phobia	13 (26%)				
Obsessive-compulsive disorder	5 (10%)				
Post-traumatic stress disorder	4 (8%)				

Note. BMI = body mass index; EDE-Q = Eating Disorder Examination-Questionnaire; FAD = McMaster Family Assessment Device; IBW = ideal body weight; STAI = Spielberger State Trait Anxiety Inventory.

Paired sample t tests were used to assess statistical significance for within-subject differences in continuous variables over time. Lower scores reflect decreased symptoms (EDE-Q, STAI) and improved family function (FAD). The EDE-Q was modified for the post-treatment evaluation to assess ED symptoms over the past 7 days. The Mini International Neuropsychiatric Interview (MINI) (Sheehan et al., 1998), a structured clinical interview designed to assess the presence of DSM-IV psychiatric disorders in adults, was used to assess current comorbid psychiatric diagnoses.

outcomes must be interpreted with caution. There was no experimental control of treatment between post-treatment and follow-up assessment, and service utilization data were not collected, rendering it difficult to attribute follow-up benefits solely to this 5-day treatment. Another significant limitation is the relatively poor follow-up assessment compliance. However, the difference in BMI from post-treatment to follow-up indicates that patients continued to gain weight, which may be attributable to continued application of skills learned during treatment or to responsiveness to other interventions after the 5-day treatment. Notably, this 5-day treatment may increase effectiveness of subsequent treatment by serving as a primer and motivating patients and their SU to engage in longer-term care. Moreover, our finding that individuals lost to follow-up were less chronic leads to the notion that this may be a promising treatment for long-standing AN. Fully-powered randomized controlled trials with longer follow-up periods will be needed to firmly establish the efficacy of this treatment.

5 | CONCLUSION

This preliminary study investigated an innovative SU-involved and neurobiologically-guided program delivered via an intense 5-day

multifamily format. Findings suggest this is a feasible, acceptable and potentially effective approach to reduce AN symptoms. This study begins to address the critical need to develop and test neurobiologically-targeted treatments for adults with AN.

ACKNOWLEDGMENTS

Support provided by the National Eating Disorders Association (NEDA) to CEW and LH, the Global Foundation for Eating Disorders (GFED) and the Price Foundation.

ORCID

Christina E. Wierenga  <http://orcid.org/0000-0002-4843-1809>

Walter H. Kaye  <http://orcid.org/0000-0002-4478-4906>

REFERENCES

- Abramowitz, J., Foa, E., & Franklin, M. (2003). Exposure and ritual prevention for obsessive-compulsive disorder: Effects of intensive versus twice-weekly sessions. *Journal of Consulting & Clinical Psychology, 71* (2), 394-398. <https://doi.org/10.1037/0022-006X.71.2.394>
- American Psychiatric Association (2000). *Diagnostic and statistical manual of mental disorders: DSM-VI-TR* (4th ed.). Washington, DC: Association AP.

- Bardone-Cone, A. M., Higgins, M. K., St. George, S. M., Rosenzweig, I., Schaefer, L. M., Fitzsimmons-Craft, E. E., . . . Preston, B. F. (2016). Behavioral and psychological aspects of exercise across stages of eating disorder recovery. *Eating Disorders*, 24(5), 424–439. <https://doi.org/10.1080/10640266.2016.1207452>
- Berner, L., Simmons, A., Wierenga, C., Bischoff-Grethe, A., Paulus, M., & Bailer, U. (2017). Altered interoceptive activation before, during, and after aversive breathing load in women remitted from anorexia nervosa. *Psychological Medicine*, 17, 1–13. <https://doi.org/10.1017/S0033291717001635>
- Byrne, S., Wade, T., Hay, P., Touyz, S., Fairburn, C., & Treasure, J. (2017). A randomised controlled trial of three psychological treatments for anorexia nervosa. *Psychological Medicine*, 29, 1–11. <https://doi.org/10.1017/S0033291717001349>
- Dare, C., Eisler, I., Russell, G., Treasure, J., & Dodge, L. (2001). Psychological therapies for adults with anorexia nervosa. Randomised controlled trial of out-patient treatments. *British Journal of Psychiatry*, 178, 216–221.
- DeGuzman, M., Shott, M., Yang, T., Riederer, J., & Frank, G. (2017). Association of elevated reward prediction error response with weight gain in adolescent anorexia nervosa. *American Journal of Psychiatry*, 174(6), 557–565. <https://doi.org/10.1176/appi.ajp.2016.16060671>
- Dimitropoulos, G., Farquhar, J., Freeman, V., Colton, P., & Olmsted, M. (2015). Pilot study comparing multi-family therapy to single family therapy for adults with anorexia nervosa in an intensive eating disorder program. *European Eating Disorders Review*, 23(4), 294–303. <https://doi.org/10.1002/erv.2359>
- Eisler, I., L., Grange, D., & Lock, J. (2015). Family therapy for adolescent eating disorders. In T. Sexton & J. Lebow (Eds.), *Handbook of family therapy* (4th ed.). Routledge. <https://doi.org/ISBN: 1138917621>
- Eisler, I., Simic, M., Hodsoll, J., Asen, E., Berelowitz, M., Connan, F., . . . Landau, S. (2016). A pragmatic randomised multi-centre trial of multi-family and single family therapy for adolescent anorexia nervosa. *BMC Psychiatry*, 16(1), 422. <https://doi.org/10.1186/s12888-016-1129-6>
- Epstein, N., Baldwin, L., & Bishop, D. (1983). The McMaster family assessment device. *Journal of Marital & Family Therapy*, 9(2), 171–180. <https://doi.org/10.1080/00224540209603911>
- Fairburn, C. G., & Beglin, S. (1992). Assessment of eating disorders: Interview or self-report questionnaire? *International Journal of Eating Disorders*, 12(1), 113–370. [https://doi.org/10.1002/1098-108X\(199412\)16:4<363::AID-EAT2260160405>3.0.CO;2-#](https://doi.org/10.1002/1098-108X(199412)16:4<363::AID-EAT2260160405>3.0.CO;2-#)
- Fassino, S., Piero, A., Gramaglia, C., & Abbate-Daga, G. (2004). Clinical, psychopathological and personality correlates of interoceptive awareness in anorexia nervosa, bulimia nervosa and obesity. *Psychopathology*, 37(4), 168–174. <https://doi.org/10.1159/000079420>
- First, M. B., Gibbon, M., Spitzer, R. L., & Williams, J. B. W. (1996). *Users guide for the structured clinical interview for DSM-IV Axis I disorders—Research version (SCID-I, version 2.0, February 1996 Final Version)*. New York: Biometrics Research Department, New York State Psychiatric Institute.
- Gallo, K. P., Chan, P. T., Buzzella, B. A., Whitton, S. W., & Pincus, D. B. (2012). The impact of an eight-day intensive treatment for adolescent panic disorder and agoraphobia on comorbid diagnosis. *Behavioural Therapy*, 43(1), 153–159. <https://doi.org/10.1016/j.beth.2011.05.002>
- Harrison, A., O'Brien, N., Lopez, C., & Treasure, J. (2010). Sensitivity to reward and punishment in eating disorders. *Psychological Research*, 177(1–2), 1–11. <https://doi.org/10.1016/j.psychres.2009.06.010>
- Hay, P., Touyz, S., & Sud, R. (2012). Treatment for severe and enduring anorexia nervosa: A review. *Australian & New Zealand Journal of Psychiatry*, 46(12), 1136–1144. <https://doi.org/10.1177/0004867412450469>
- Hill, L. (2017). Can your brain cure anorexia? A brain-based approach to eating disorder treatment eating disorders catalogue. <https://doi.org/www.edcatalogue.com/hill/#more-4635>
- Hill, L., Peck, S., Wierenga, C., & Kaye, W. (2016). Applying neurobiology to the treatment of adults with anorexia nervosa. *Journal of Eating Disorders*, 4(1), 31. <https://doi.org/10.1186/s40337-016-0119-x>
- Kaye, W. H., Wierenga, C. E., Knatz, S., Liang, J., Boutelle, K., Hill, L., & Eisler, I. (2015). Temperament-based treatment for anorexia nervosa. *European Eating Disorders Review*, 23(1), 12–18. <https://doi.org/10.1002/erv.2330>
- Kerr, K., Moseman, S., Avery, J., Bodurka, J., Zucker, N., & Kyle Simmons, W. (2016). Altered insula activity during visceral interoception in weight-restored patients with anorexia nervosa. *Neuropsychopharmacology*, 41(2), 521–528. <https://doi.org/10.1038/npp.2015.174>
- Knatz, S., Wierenga, C., Murray, S., Hill, L., & Kaye, W. (2015). Neurobiologically-informed treatment for adults with anorexia nervosa: A novel approach to a chronic disorder. *Dialogues in Clinical Neuroscience*, 17(2), 229–236.
- Lemmens, G., Eisler, I., Buysse, A., Heene, E., & Demyttenaere, K. (2009). The effects on mood of adjunctive single family and multi-family group therapy in the treatment of hospitalised patients with major depression: An RCT and 15 months follow-up study. *Psychotherapy & Psychosomatics*, 78(2), 98–105. <https://doi.org/10.1159/000201935>
- Lilenfeld, L. (2011). Personality and temperament. *Current Topics in Behavioural Neuroscience*, 6, 3–9. https://doi.org/10.1007/7854_2010_86
- Lock, J. (2015). An update on evidence-based psychosocial treatments for eating disorders in children and adolescents. *Journal of Clinical Child & Adolescent Psychology*, 44(5), 707. <https://doi.org/10.1080/15374416.2014.971458>
- Loeb, K. (2008). Eating Disorder Examination Questionnaire—Parent Version (P-EDEQ), Version 1.4. Unpublished measure based on Fairburn, C.G. & Beglin, S.J. The Eating Disorder Examination Questionnaire (EDE-Q). In C. G. Fairburn (Ed.), *Cognitive behavior therapy and eating disorders*. New York: Guilford Press.
- Marzola, E., Knatz, S., Murray, S. B., Rockwell, R., Boutelle, K., Eisler, I., & Kaye, W. H. (2015). Short-term intensive family therapy for adolescent eating disorders: 30-month outcome. *European Eating Disorders Review*, 23(3), 210–218. <https://doi.org/10.1002/erv.2353>
- McFarlane, W. (2016). Family interventions for schizophrenia and the psychoses: A review. *Family Process*, 55(3), 460–482. <https://doi.org/10.1111/famp.12235>
- Miller, I. W., Solomon, D. A., Ryan, C. E., & Keitner, G. (2004). Does adjunctive family therapy enhance recovery from bipolar I mood episodes? *Journal of Affective Disorders*, 82(3), 431–436. <https://doi.org/10.1016/j.jad.2004.01.010>
- NICE (2017). *Eating disorders: Recognition and treatment. NICE guideline [NG69]*. National Institute for Health Care.
- Oberndorfer, T., Frank, G., Fudge, J., Simmons, A., Paulus, M., & Wagner, A. (2013). Altered insula response to sweet taste processing after recovery from anorexia and bulimia nervosa. *American Journal of Psychiatry*, 170(2), 132–141. <https://doi.org/10.1176/appi.ajp.2013.111.11745>
- Rockwell, R., Boutelle, K., Trunko, M., Jacobs, M., & Kaye, W. (2011). An innovative short-term, intensive, family-based treatment for adolescent anorexia nervosa: Case series. *European Eating Disorders Review*, 19(4), 362–367. <https://doi.org/10.1002/erv.1094>

- Schmidt, U., & Campbell, I. (2013). Treatment of eating disorders cannot remain 'brainless': The case for brain-directed treatments. *European Eating Disorders Review*, 21(6), 425–427. <https://doi.org/10.1002/erv.2257>
- Schmidt, U., Renwick, B., Lose, A., Kenyon, M., DeJong, H., Broadbent, H., ... Landau, S. (2013). The MOSAIC study—Comparison of the Maudsley Model of Treatment for Adults with Anorexia Nervosa (MANTRA) with Specialist Supportive Clinical Management (SSCM) in outpatients with anorexia nervosa or eating disorder not otherwise specified, anorexia nervosa type: Study protocol for a randomized controlled trial. *Trials*, 14(1), 160. <https://doi.org/10.1186/1745-6215-14-160>.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* 59(20), 22–33.
- Spielberger, C., Gorsuch, R., & Lushene, R. (1970). *STAI manual for the state trait anxiety inventory*. Palo Alto, CA: Consulting Psychologists Press.
- Steinhausen, H. C. (2002). The outcome of anorexia nervosa in the 20th century. *Am J Psychiatry*, 159(8), 1284–1293. 12153817
- Storch, E. A., Geffken, G. R., Merlo, L. J., Mann, G., Duke, D., Munson, M., ... Goodman, W. K. (2007). Family-based cognitive-behavioral therapy for pediatric obsessive-compulsive disorder: Comparison of intensive and weekly approaches. *Journal of American Academy of Child & Adolescent Psychiatry*, 46(4), 469–478. <https://doi.org/https://doi.org/10.1097/chi.0b013e31803062e7>
- Tantillo, M. (2006). A relational approach to eating disorders multifamily therapy group: Moving from difference and disconnection to mutual connection. *Families, Systems, & Health*, 24(1), 82–102. <https://doi.org/10.1037/1091-7527.24.1.82>
- Wagner, A., Barbarich-Marsteller, N. C., Frank, G. K., Bailer, U. F., Wonderlich, S. A., Crosby, R. D., ... Kaye, W. H. (2006). Personality traits after recovery from eating disorders: Do subtypes differ? *International Journal of Eating Disorders*, 39(4), 276–284. <https://doi.org/10.1002/eat.20251>
- Watson, H., & Bulik, C. (2013). Update on the treatment of anorexia nervosa: Review of clinical trials, practice guidelines and emerging interventions. *Psychological Medicine*, 43(12), 2477–2500. <https://doi.org/10.1017/S0033291712002620>
- Wierenga, C. E., Bischoff-Grethe, A., Melrose, A. J., Irvine, Z., Torres, L., Bailer, U. F., ... Kaye, W. H. (2015). Hunger does not motivate reward in women remitted from anorexia nervosa. *Biological Psychiatry*, 77(7), 642–652. <https://doi.org/10.1016/j.biopsych.2014.09.024>

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Wierenga CE, Hill L, Knatz Peck S, et al. The acceptability, feasibility, and possible benefits of a neurobiologically-informed 5-day multifamily treatment for adults with anorexia nervosa. *Int J Eat Disord*. 2018;00:1–7. <https://doi.org/10.1002/eat.22876>